



2012 International Conference on Solid State Devices and Materials Science

## Rapid Determination Method for 12 Pyrethroid Pesticide Residues in Tea by Stir Bar Sorptive Extraction-Thermal Desorption-Gas Chromatography\*

Bin Li, Fangang Zeng, Qichun Dong, Yang Cao, Haitao Fan, Chuanfeng Deng

*School of Environment and Natural Resources, Renmin University of China, Beijing 100872, China*

### Abstract

The 12 pyrethroid Pesticides residues in tea samples can be rapidly extracted and separated by a novel stir bar sorptive extraction (SBSE)-thermal desorption (TDU)-gas chromatography (GC) method. Impurities and pyrethroid pesticides residues of tea can be separated by this method, and then use TDU-GC to analyse pyrethroid pesticides residues. The residues of 12 pyrethroid pesticides at the same time can be identified and quantified simultaneously by SBSE-TDU-GC rapidly with high sensitivity and good reproducibility. This method is simple, rapid, and the average adding standard recovery ratios were 92.89%-105.01%. The correlation coefficient of pyrethroid pesticides in tea samples is  $\geq 0.9926$ . Good repeatability ( $n=6$ ) was obtained in all the cases with the relative standard deviation lower 9.7%. Pyrethroid pesticides residues in 4 real tea samples can be determined rapidly by this method, their pyrethroid pesticide residues are all less than MRL of China and European Union.

© 2012 Published by Elsevier B.V. Selection and/or peer-review under responsibility of Garry Lee

Open access under [CC BY-NC-ND license](http://creativecommons.org/licenses/by-nc-nd/4.0/).

**Keywords:** pyrethroid pesticide; tea; stir bar sorptive extraction (SBSE); thermal desorption; GC-ECD

European Union issued a number of maximum residue limits (MRL) of pesticides in food standard on August 27th 2004. This standard specifies 168 MRL criterion of pesticides in tea items, including some pyrethroid pesticides. Pyrethroids are a class of synthetic esters, which are derived by changing the chemical structure of natural pyrethrins. Its insecticidal toxicity is increased by 10 to 100 times than the older generation of pesticides (1). Pyrethroids are toxic organic pollutants, which have caused widespread concern in the international, and some of them have carcinogenic, teratogenic and mutagenic effects. The traditional analysis procedure for pyrethroid pesticide residues in tea has some problems, such as complicated sample pretreatment, using a large number of organic solvents and so on (2-5). To this end, it is very necessary to work up a multi-residue analytical method with high efficiency, high sensitivity and strong selectivity. In this paper, the pyrethroid pesticide residues in tea would be determined by stir bar sorptive extraction (SBSE) method. Stir bar sorptive extraction (SBSE) is constituted by the stir bar coated with Poly-dimethyl-siloxane (PDMS) (6). The principles of SBSE and solid phase micro extraction (SPME) are same, but the

## 1. Materials and Methods

This work is supported by the Asian Research Center of Renmin University of China

## 2. Results and Analysis

### 2.1 Influence Factors of the Extraction Efficiency

#### 2.1.1 Extraction Time

Under the same experimental conditions (certain rotor speed and extraction temperature), adding 4  $\mu\text{L}$  mixed standard solution (fenson, allethrin, Ovex, tetramethrin, fenpropathrin, permethrin, cypermethrin, deltamethrin, fenvalerate, bifenthrin, cyfluthrin are in the concentration of  $2\mu\text{g/L}$ ) into six 20 mL headspace bottles which were added with 20 mL pure water already. Then in different extraction time (0.25, 0.5, 1, 2, 4, 8 h), comparing the extraction efficiency according to the chromatographic peak area (peak area of each pyrethroid is the sum of all of its isomers' peak area) of 12 pyrethroids. The results indicated that the total peak area of 12 pyrethroids were increased with increase of the extraction time. The extraction was not reached equilibrium in 8 h, but when the extraction time was 1 h, peak area growth rate slowed down. And the peak area got a fastest growing in the range of 0 - 1 h. In order to achieve the purpose of rapid determination, the best extraction time was designated as 1h.

#### 2.1.2 The Influence of Methanol on Extraction Efficiency

As the tea samples is solid, it can not be extracted directly. So before the sorption extraction, methanol extraction was implemented, and then the samples of methanol extraction was extracted by stir bar sorptive extraction (SBSE) method after diluted by distilled water. Methanol in extraction diluted with distilled water may have effects on the coefficient of recovery of Aimed Compound, so methanol extraction was replaced with methanol to investigate the effects of volume fraction of methanol (0, 10%, 20%, 40% and 60%) on extraction volume. The results showed that the methanol from 0-20% can strengthen stir bar to target extraction. But when the volume fraction range from 40% to 60%, stir bar to target extraction of Aimed Compound was weakened. The reason may be that the additional methanol reduce the adsorption of compound by Vial wall, but the High concentrations of methanol make PDMS have effect on The adsorption of compound. When volume fraction of methanol was added from 0 to 10%, Peak area almost Compounds was augmented largely. In order to save solvent, the methanol whose volume fraction was 10% has been used in this experiment. That is to say, 2 mL of methanol extraction was diluted with distilled water before Pretreatment with SBSE.

### 2.2 Optimization of Conditions of Thermal Desorption

At the same conditions of SBSE, Desorption temperature, Desorption time and CIS-4 inlet temperature of cold trap influence the thermal desorption Aimed Compound from Stirring rod and entrance into Capillary column.

#### 2.2.1 Optimization of Desorption Time

With the desorption time range from 20 to 40min, the total peak area was increased, which was because that The longer thermal desorption time lead the More volatilization of organic on Stirring rod. in view of the slow increase of peak area when the desorption time range from 20 to 40min, 30min of desorption time was used to control the Experimental time in this Experiment.

#### 2.2.2 Optimization of Desorption Temperature

With the desorption temperature range from 250 to 300°C, the total peak area of all pyrethroids were increased, which was because that the higher desorption temperature lead more volatilization of organic compound. 300°C desorption temperature was used in this Experiment.

#### 2.2.3 Optimization of CIS-4 Inlet Temperature of Cold Trap

With the CIS-4 inlet temperature range from 0°C to below zero, The total peak area was reduced in turn. 0°C of CIS-4 inlet temperature was used in this experiment.

The results showed that 300°C desorption temperature, 30min of desorption time and 0°C inlet temperature of cold trap were the best conditions.

### 2.3 Evaluation of Method

Using standard addition method, use blank sample added with guide sample to quantify, to eliminate the of sample matrix.in this experiment, Mixed standard solution whose concentration range from 0.3-3 µg/ml and 8 ml of methanol were added to 2 g blank tea sample after ultrasonic extraction for 30 min, 2ml of Upper liquid was moved to 20 ml vial added by 18 ml of purified water for Stir bar sorptive extraction. The extraction time and methanol concentration were 1h and 10%. The experiment was been repeated for 6 times to Measure peak area and make standard curve under the best conditions. Correlation coefficient, relative standard deviation and recoveries were showed in Table 1. Analysis results meet the requirements of pesticide residues analysis.

Table 1 Correlation coefficient ,relative standard deviation, recoveries

Pyrethroid	Linea- rities	RSD (n=6,%)	Recoveries (n=6,%)
Fenson	0.9935	5.2	95.37
Allethrin	0.9926	9.7	92.89
Ovex	0.9928	5.1	105.01
Tetramethrin Fenpropathrin	0.9991	5.1	101.13
Permethrin	0.9942	5.5	96.57
Trans-cypermethrin	0.9937	5.6	104.06
Cis-cypermethrin	0.9969	7.6	101.16
Deltamethrin	0.9986	7.6	99.26
Fenvalerate	0.9953	9.5	97.06
Bifenthrin	0.9970	3.6	97.20
Cyfluthrin	0.9996	5.1	104.48
	0.9980	3.0	101.04

### 2.4 Analysis of Real Samples

4 tea samples purchased at random were detected for 3 Parallels using the above methods. Average were showed in Table 2. The results show that pyrethroid residues in tea samples did not exceed the stringent limits made by China and the EU(14-15).

Table 2 Pyrethroid Pesticides in Tea Samples (µg • kg<sup>-1</sup>)

Pyrethroid	Pu'er Tea	DaFang Tea	Jasmine Tea	Jasmine Beads Tea
Fenson	—	0.0798	—	—
Allethrin	0.0403	—	—	—
Ovex	—	—	—	—
Tetramethrin	—	—	0.0901	—
Fenpropathrin	—	—	—	0.05243
Permethrin	—	—	0.0806	—
Trans-cypermethrin	—	0.0975	—	—
Cis-cypermethrin	—	—	—	0.5383
Deltamethrin	—	—	—	—
Fenvalerate	0.0409	—	—	—
Bifenthrin	—	—	—	—
Cyfluthrin	—	—	—	—

Note: — means less than detection limit.

### 3. Conclusion

This study established a new method which use SBSE-TDU-GC to detect the pyrethroid pesticides in tea, and the correlation coefficient( $\geq 0.9926$ ), The relative standard deviation( $\leq 9.7$ ,  $n=6$ ) and recovery ratio(92.89%-105.01%) reached the testing requirements of pyrethroid in tea. The method have some virtues such as relatively simple pretreatment, short processing time and good repeatability. Besides, the method was applicable to pyrethroid pesticide residue detection requirements of tea. SBSE-TDU-GC rapid determination may be used in detection of other food pesticide widely in the future. on the other hand, in order to get optimum results of specific components in specific sample, More appropriate methods and testing conditions should be established.

As can be seen from the experimental results, Cypermethrin, permethrin and tetramethrin residues detected in 4 tea samples were relatively more, which means that the use of these types of pesticides in teaplantation should be noticed. Pyrethroid residues in tea samples of China did not exceed the stringent limits made by China and the EU.

### References

- [1] Jieping Cao. Brief content of the issue of pesticide residues in tea exports. *Journal of Tea Industry*, vol. 3, no.21, pp. 23-24, 1999.
- [2] Lili Yang, Ying Ji, Yuanping He. Determination of pyrethrins residues in tea by gas chromatography. *Chemical Analysis and Meterage*, vol.5, no.14, pp.38-40, 2005.
- [3] Huiying Ding; Xiaoxia Bao; Zheng Ziqiang. Determination of 7 pyrethroids residues in tea. *Journal of Instrumental Analysis*, vol.6, no.19, pp.31-34, 2000.
- [4] Zhong Liu, Kejin Xie, Qixiang Chen and Shenghua Wang. GC determination of remains of seven kinds of pyrethroid pesticide in tea samples. *Analytical Laboratory*, vol.6, no.21, pp.103-104, 2002
- [5] Ying Zhang, Zhiqiang Huang, Yongjun Li, et al. Gas chromatography in Tea pyrethroid pesticide residues. *Chinese Journal of Analytical Chemistry*, vol.3, no.30, pp.377, 2002.
- [6] Baltussen E, Sandra P, David F, Cramers C.J. A novel extraction technique for aqueous samples: theory and principles. *J Microcol Sep.*, 1999, pp.737.
- [7] Baltussen E, Tienpont B, David F. Multi-residue screening of pesticides in vegetables, fruits and baby food by stir bar sorptive extraction-thermal desorption-capillary gas chromatography-mass spectrometry. *J Chromatogr A*, vol.1000, pp.299, 2003.
- [8] Wenmin Liu, Yuan Hu, Jing-hong Zhao, et al. Determination of organophosphorus pesticides in cucumber and potato by stir bar sorptive extraction. *J Chromatogr A*, 2005, vol.1095, pp.1.
- [9] Juan Garcia A, Pico Y, Font G. Capillary electrophoresis for analyzing pesticides in fruits and vegetables using solid-phase extraction and stir-bar sorptive extraction. *J Chromatogr A*, 2005, vol. 1073, pp.229.
- [10] Blasco C, Font G, Pico Y. Comparison of microextraction procedures to determine pesticides in oranges by liquid chromatography-mass spectrometry. *J Chromatogr A*, 2002, vol.970, pp. 201.
- [11] Blasco C, Frenandez M, Pico Y, et al. Comparison of solid phase microextraction and stir bar sorptive extraction for determining six organophosphorus insecticides in honey by liquid chromatography-mass spectrometry. *J Chromatogr A*, 2004, vol.1030, pp.77.
- [12] Yuan Xu, Wenmin Liu, Jinghong Zhao, Yafeng Guan. Determination of polycyclic aromatic hydrocarbons in seawater using stir bar sorptive extraction-gas chromatography. *Chinese Journal of Analytical Chemistry*, vol.10, no.33, pp. 1401, 2005.
- [13] Wenmin Liu, Yuan Xu, Wenna Guan, Yafeng Guan. Solid Phase microextraction(SPME)technology- stir bar sorptive extraction (SBSE). *Modern Scientific Instruments*, vol.1, pp.51-53, 2006.
- [14] Zongmao Chen. New standards for pesticide residues in tea issued by European Union and Japan . *China's Tea Industry*, vol.6, pp.12-13, 2004.
- [15] GB2763—2005 Maximum residue limits for pesticides in food[M], Beijing: Standards Press of China, 2005.